

STOKES THEOREM FOR LIPSCHITZ FORMS ON A SMOOTH MANIFOLD

STANISLAV DUBROVSKIY

ABSTRACT. Stokes theorem holds for Lipschitz forms on a smooth manifold.

Let ω be a Lipschitz $(n-1)$ -form with compact support K on a smooth n -manifold M , in the notation of [GKS]:

$$\omega \in \mathcal{L}^{n-1}(M) .$$

Suppose first $M = \mathbb{R}^n$ or $\mathbb{H}^n = \{x_n \geq 0\}$, the upper half space in $\mathbb{R}^n$. It suffices to assume

$$\omega = f dg_1 \wedge dg_2 \dots \wedge dg_{n-1}$$

(rather than a finite sum of such terms), where $f, g_1, g_2, \dots, g_{n-1}$ are Lipschitz functions on M . According to [GKS, Lemma 1.1]:

$$d\omega = df \wedge dg_1 \wedge dg_2 \dots \wedge dg_{n-1} .$$

We can write

$$\omega = a dx_1 \wedge dx_2 \dots \wedge dx_{n-1} + \omega_n ,$$

where ω_n contains dx_n , and a is a polynomial in $f, \frac{\partial g_i}{\partial x_j}$.

Then

$$d\omega = b dx_1 \wedge dx_2 \dots \wedge dx_n ,$$

b is a polynomial in $\frac{\partial f_i}{\partial x_j}, \frac{\partial g_i}{\partial x_j}$.

Consider $a|_{x_n=0}$, the restriction of a to $K' = K \cap \{x_n = 0\}$, a compact subset in K . It is again a polynomial in Lipschitz functions, since restrictions of Lipschitz functions to a hyperplane are Lipschitz.

Just as in the proof of [GKS, Lemma 1.1] let us consider smooth forms

$$\omega_s = f_{\varepsilon_s} d(g_1)_{\varepsilon_s} \wedge d(g_2)_{\varepsilon_s} \dots \wedge d(g_{n-1})_{\varepsilon_s}$$

and

$$d\omega_s = df_{\varepsilon_s} d(g_1)_{\varepsilon_s} \wedge d(g_2)_{\varepsilon_s} \dots \wedge d(g_{n-1})_{\varepsilon_s}$$

- sequence ε_s chosen so that the Steklov averages $f_{\varepsilon_s}, (g_1)_{\varepsilon_s}, (g_2)_{\varepsilon_s}, \dots, (g_{n-1})_{\varepsilon_s}$ have their derivatives converge:

$$\left(\frac{\partial f}{\partial x_j} \right)_{\varepsilon_s} \rightarrow \frac{\partial f}{\partial x_j} , \quad \left(\frac{\partial g_i}{\partial x_j} \right)_{\varepsilon_s} \rightarrow \frac{\partial g_i}{\partial x_j}$$

almost everywhere on K and K' .

(This can be done since the restrictions to K' are still Lipschitz functions.)

Date: May 27, 2008.

Key words and phrases. Lipschitz forms, Stokes theorem.

Functions

$$a(f_{\varepsilon_s}, (\frac{\partial g_i}{\partial x_j})_{\varepsilon_s}) \quad , \quad b((\frac{\partial f_i}{\partial x_j})_{\varepsilon_s}, (\frac{\partial g_i}{\partial x_j})_{\varepsilon_s})$$

are uniformly bounded in s and converge almost everywhere to

$$a(f, \frac{\partial g_i}{\partial x_j}) \quad , \quad b(\frac{\partial f_i}{\partial x_j}, \frac{\partial g_i}{\partial x_j})$$

respectively.

Then by the Lebesgue limit theorem:

$$\int_{\partial M} \omega_s \rightarrow \int_{\partial M} \omega \quad , \quad \int_M d(\omega_s) \rightarrow \int_M d\omega .$$

Since Stokes theorem is valid for smooth ω_s , we must conclude it holds for the Lipschitz forms in the limit.

Suppose now M consists of a single chart $\phi : M \rightarrow \mathbb{R}^n$ (or $\mathbb{H}^n$).

We have:

$$\int_{\partial M} \omega = \int_{\partial(\phi(M))} (\phi^{-1})^* \omega = \int_{\phi(M)} (\phi^{-1})^* d\omega = \int_M d\omega .$$

For an atlas with multiple charts, we use partition of unity $\{\rho_\alpha\}$ and the fact that $\rho_\alpha \omega$ are again Lipschitz, together with the linearity of integral, cf. [BT].

Integration on Lipschitz manifolds is to be discussed in a forthcoming paper.

REFERENCES

- [BT] Bott R., Tu L., Differential forms in Algebraic Topology. Graduate Texts in Mathematics, **82**, Springer-Verlag, New York-Berlin, 1982.
- [GKS] Gol'dshtein V.M., Kuz'minov V.I., Shvedov I.A., Differential forms on Lipschitz Manifolds, *Siberian Math. Journal*, **23**(2)(1982), 16-30.

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, P.O.B. 653, BEER'SHEVA
84105, ISRAEL

E-mail address: `dubr@math.bgu.ac.il`